arXiv:1609.00595v1 [math.AC] 2 Sep 2016

ON THE RELATIONS OF ISOTONIAN ALGEBRAS

JÜRGEN HERZOG, AYESHA ASLOOB QURESHI AND AKIHIRO SHIKAMA

Abstract. It is shown that for large classes of posets P and Q, the defining ideal $J_{P,Q}$ of an isotonian algebras is generated by squarefree binomials. Within these classes, those posets are classified for which $J_{P,Q}$ is quadratically generated.

Introduction

In [1], isotonian algebras were introduced as a generalization of Hibi rings. Given two finite posets P and Q, the algebra $K[P,Q]$ is generated by monomials

$$u_\varphi = \prod_{p\in P} x_{p,\varphi(p)} \quad \text{with} \quad p \in P,$$

where $\varphi : P \to Q$ is an isotone (order preserving) map. We denote the defining ideal of $K[P,Q]$ by $J_{P,Q}$. Since $K[P,Q]$ is a toric ring, the ideal $J_{P,Q}$ is generated by binomials.

Isotone maps first appeared in Commutative Algebra in the study of the ideals $I(P,Q)$ which are generated by all the monomials u_φ, see [2]. The structure of these ideals has been further studied in subsequent papers [8], [5], [9],[10], [11].

In the case that Q is a chain of length 2, the algebra $K[P,Q]$ coincides with the so-called and well studied Hibi ring, see [6], [4], [7], [3]. It is known that the defining ideals of Hibi rings admit quadratic Gröbner basis and they are normal Cohen-Macaulay domains. In the above mentioned paper [1], it is also shown that $J_{P,Q}$ admits a quadratic Gröbner basis if P is a chain and Q is a co-rooted poset. However, easy examples show that isotonian algebras need not to admit quadratic Gröbner bases. On the other hand, it is conjectured that the ideal $J_{P,Q}$ has a squarefree initial ideal for a suitable term order.

We call $f = u - v$ squarefree, if u and v are squarefree monomials. In support of the above conjecture we prove in Theorem 3.3, that $J_{P,Q}$ is generated by squarefree binomials in the case that each connected component of P is rooted or co-rooted, and that $J_{P,Q}$ is generated by quadratic binomials if and only if every poset cycle of length ≥ 6 in Q admits a chord.

1. Reduction to the case that P is connected

Let P be a finite poset. We say that P is connected if its Hasse diagram is connected as a graph. Given two posets P_1 and P_2, the sum $P_1 + P_2$ is defined to be the disjoint union of the elements of P_1 and P_2 with $p \leq q$ if and ony if $p, q \in P_1$

2010 *Mathematics Subject Classification.* 05E45, 05E40, 13C99.

This paper was partially written during the stay of the second author at The Abdus Salam International Centre of Theoretical Physics (ICTP), Trieste, Italy.

or $p, q \in P_2$ and $p \leq q$ in the corresponding posets P_1 or P_2. Then its is clear that P can be written as $P = P_1 + P_2 + \cdots + P_r$ where each P_i is connected. The P_i are called the connected components of P.

Let $P, P_1, P_2, \ldots, P_r$ and $Q, Q_1, Q_2, \ldots, Q_s$ be finite posets. Then it is shown in [1, Lemma 1.3] that $K[\sum_{i=1}^r P_i, Q]$ is isomorphic to the Segre product $K[P_1, Q] * K[P_2, Q] * \cdots * K[P_r, Q]$. Recall that if $R_1 = K[f_1, \ldots, f_r]$ and $R_2 = K[g_1, \ldots, g_s]$ are two standard graded K-algebras. Then

$$R_1 * R_2 = K[\{f_i g_j \colon\ i = 1, \ldots, r, j = 1, \ldots, s\}]$$

is the Segre product of R_1 and R_2.

We denote by $\mathrm{Hom}(P, Q)$ the set of isotone maps from $P \to Q$, and let $T = K[t_\varphi \colon\ \varphi \in \mathrm{Hom}(P, Q)]$ be the polynomial ring in the variables t_φ, and let $\varphi : T \to K[P, Q]$ be the K-algebra homomorphism defined by $t_\varphi \mapsto \prod_{p \in P} x_{p\varphi(p)}$. We denote the kernel of φ by $I_{P,Q}$.

Theorem 1.1. *Let P and Q be two finite posets and let $P_1, P_2, \ldots, P_r$ be the connected components of P. Then $I_{P,Q}$ is generated by squarefree, respectively quadratic binomials, if each $I_{P_i,Q}$ is generated by squarefree, respectively quadratic binomials.*

Proof. By using induction on r it suffices to prove the following fact: Let $R_1 = K[f_1, \ldots, f_r]$ and $R_2 = K[g_1, \ldots, g_s]$ be standard graded toric rings whose defining ideals are generated by squarefree binomials, then this is also the case for $R_1 * R_2$. To show this we introduce some notation: let $T_1 = K[x_1, \ldots, x_r]$ and $T_2 = K[y_1, \ldots, y_s]$ be the polynomial ring over K, and I_1 the kernel of K-algebra homomorphism $T_1 \to R_1$ with $x_i \mapsto f_i$. The ideal I_2 is similarly defined. Next, let $T = K[z_{ij} \mid i = 1, \ldots, r, j = 1, \ldots, s]$ be the polynomial ring over K and I the kernel of the K-algebra homomorphism $T \to R_1 * R_2$ with $z_{ij} \mapsto f_i g_j$.

We have to show that any binomials $h = \prod_{t=1}^d z_{i_t j_t} - \prod_{t=1}^d z_{k_t l_t} \in I$ is a linear combination of squarefree, respectively quadratic binomials of I. Note that $h = h_1 + h_2$, where

$$h_1 = \prod_{t=1}^d z_{i_t j_t} - \prod_{t=1}^d z_{k_t j_t}, \quad h_2 = \prod_{t=1}^d z_{k_t j_t} - \prod_{t=1}^d z_{k_t l_t}.$$

Observe that $h_1, h_2 \in I$. Indeed, $h_1 \mapsto \prod_{t=1}^d (f_{i_t} g_{j_t}) - \prod_{t=1}^d (f_{k_t} g_{j_t})$, and we have to show that $\prod_{t=1}^d (f_{i_t} g_{j_t}) - \prod_{t=1}^d (f_{k_t} g_{j_t}) = 0$.

Since $h \in I$, it follows that $\prod_{t=1}^d (f_{i_t} g_{j_t}) - \prod_{t=1}^d (f_{k_t} g_{l_t}) = 0$. Let $R_2 \to K$ be the K-algebra homomorphism such that $g_i \mapsto 1$. This K-algebra homomorphism induces a canonical K-algebra homomorphism $R_1 * R_2 \to R_1 * k = R_1$. This shows $\prod_{t=1}^d (f_{i_t} g_{j_t}) - \prod_{t=1}^d (f_{k_t} g_{l_t}) = 0$ is mapped to $\prod_{t=1}^d f_{i_t} - \prod_{t=1}^d f_{k_t} = 0$ in R_1. Similarly we see that $\prod_{k=1}^d g_{i_s} - \prod_{k=1}^d g_{j_k} = 0$ in R_2.

Next, let $R_1 * R_2 \to R_1 \otimes R_2 = K[f_1, \ldots f_r, g_1, \ldots, g_s]$ be the canonical inclusion map. Then, $\prod_{t=1}^d (f_{i_t} g_{j_t}) - \prod_{t=1}^d (f_{k_t} g_{j_t}) \mapsto (\prod_{t=1}^d f_{i_t} - \prod_{t=1}^d f_{k_t}) \prod_{t=1}^d g_{j_t} = 0$.

Therefore,

$$\prod_{t=1}^{d}(f_{i_t}g_{j_t}) - \prod_{t=1}^{d}(f_{k_t}g_{j_t}) = 0.$$

Similarly,

$$\prod_{t=1}^{d}(f_{k_t}g_{j_t}) - \prod_{k=1}^{d}(f_{k_t}g_{l_t}) = 0.$$

This shows that $h_1, h_2 \in I$.

Since $w = \prod_{i=1}^{d} x_{i_t} - \prod_{i=1}^{d} x_{k_t} \in I_1$, our assumption implies that w can be written as linear combination of squarefree, respectively quadratic binomials w_q. Then $w = \sum_q u_q w_q$ where u_q is a monomial in T_1, and $w_q \in I_1$ is squarefree, respectively quadratic. Let $g = \prod_{t=1}^{d} y_{j_t}$, and for each q write $g = g_{1,q}g_{2,q}$, where $g_{2,q}$ is the product of $\deg w_q$ many factors of g and $g_{1,q}$ is the product of the remaining factors. Then $w = \sum_q u_q w_q$ yields the equation

$$wg = \sum_q (u_q g_{1,q})(w_q g_{2,q}), \tag{1}$$

which is valid in $R_1 * R_2$.

Let us first assume that all w_q are squarefree binomials. Then for each q, we have $w_q = \prod_{\in V_q} x_i - \prod_{i \in V_q'} x_i$ with $V_q, V_q' \subset \{1, \ldots, r\}$ and $q_{2,q} = \prod_{j \in U_q} y_j$ with $U_q \subset \{j_1, \ldots, j_d\}$ and $|U_q| = |V_q| = |V_q'|$. For each q, let $\sigma_q : V_q \to U_q$ and $\tau_q : V_q \to U_q$ be bijections. Then (1) implies that h_1 is a linear combination of squarefree binomials of the form

$$\prod_{i \in V_q} z_{i\sigma_q(i)} - \prod_{i \in V_q'} z_{i\tau_q(i)}.$$

The similar statement holds for h_2. This shows that h is a linear combination of squarefree binomials. Similarly, if we assume all w_q are quadratic binomials, we deduce in the same way that h is a linear combination of quadratic binomials. □

2. Special binomials

In this section we study the relations of $K[P, Q]$ in the case that P is rooted. Analogous arguments hold when P is co-rooted.

To define special binomial relations, we first prove the following lemma in which a new isotone map is constructed from a given one by suitable modifications.

Lemma 2.1. *Let P and Q be two posets and assume that P is rooted. Let $\varphi \in \mathrm{Hom}(P, Q)$ and let $\psi : P \to Q$ be a map satisfying the following condition: there exists $p \in P$ such that*

(i) *$\psi(p_1) \geq \psi(p_2)$, for all $p_1 \geq p_2 \geq p$,*
(ii) *$\psi(q) = \varphi(q)$ for all $q \not\geq p$,*
(iii) *if there exists a lower neighbor q of p then $\psi(p) \geq \psi(q)$.*

Then $\psi \in \mathrm{Hom}(P, Q)$.

Proof. Let $p_1, p_2 \in P$ such that $p_1 \geq p_2$. We need to show that $\psi(p_1) \geq \psi(p_2)$. If $p_1 \geq p_2 \geq p$ or $p_1, p_2 \not\geq p$, then by using (i) and (ii), we are done. The only non trivial case is when $p_1 \geq p$ and $p_2 \not\geq p$. Since $p_1 \geq p, p_2$ and P is rooted it follows that p and p_2 are comparable. Since $p_2 \not\geq p$, we must have $p_2 < p$. In particular, p admits a lower neighbor q. Then by using (i), (ii) and (iii) we get

$$\psi(p_1) \geq \psi(p) \geq \psi(q) = \varphi(q) \geq \varphi(p_2) = \psi(p_2).$$

□

Now we are ready to define special relations. Let $\varphi_1, \ldots, \varphi_d \in \mathrm{Hom}(P, Q)$, $p \in P$ and π be a permutation on the set $\{1, \ldots, d\}$. In the case that p admits a lower neighbor $p' \in P$ then we require that π satisfies $\varphi_{\pi(i)}(p) \geq \varphi_i(p')$ for all i.

Then, by Lemma 2.1, the following maps

$$\varphi_i'(q) = \begin{cases} \varphi_{\pi(i)}(q), & \text{if } q \geq p, \\ \varphi_i(q), & \text{otherwise.} \end{cases}$$

are isotone. Consider the binomial $f = \prod_{k=1}^d t_{\varphi_i} - \prod_{k=1}^d t_{\varphi_i'}$. Then $f \in J_{P,Q}$, and $f \neq 0$ if π is not identity. More generally, if $g = \prod_{k=1}^d t_{\varphi_i} - \prod_{k=1}^d t_{\psi_i}$, Then $g \in J_{P,Q}$ if and only if for all $p \in P$

$$\{\varphi_1(p), \varphi_2(p), \ldots, \varphi_d(p)\} = \{\psi_1(p), \psi_2(p), \ldots, \psi_d(p)\} \tag{2}$$

as multi-set.

The binomial $f = \prod_{k=1}^d t_{\varphi_i} - \prod_{k=1}^d t_{\varphi_i'}$ is called *special* of type (p, π) with respect to $\varphi_1, \ldots, \varphi_d \in \mathrm{Hom}(P, Q)$. Note that f is also special of type (p, π^{-1}) with respect to $\psi_1, \ldots, \psi_d$.

Now we are ready to prove the following

Theorem 2.2. *Let P be a rooted or co-rooted poset. Then for any Q, the ideal $J_{P,Q}$ is generated by special binomials.*

Proof. Let P be a rooted poset and p_0 be the minimal element of P. Also let

$$f = t_{\varphi_1} t_{\varphi_2} \cdots t_{\varphi_d} - t_{\psi_1} t_{\psi_2} \cdots t_{\psi_d}$$

be a nonzero binomial in $J_{P,Q}$. After relabelling of ψ_i, we may assume that $\varphi_i(p_0) = \psi_i(p_0)$ for all i. Let $\mathcal{I} \subset P$ be the largest poset ideal with the property that $\varphi_i(p) = \psi_i(p)$ for all $p \in \mathcal{I}$. Since $f \neq 0$, it implies that $\mathcal{I} \neq P$.

Let $p' \in P \setminus \mathcal{I}$ such that $\mathcal{I}' = \mathcal{I} \cup \{p'\}$ is again the poset ideal in P. Note that if p' has a lower neighbor then it belongs to $\mathcal{I}$.

We will show that $f = g + h$ where g is a special binomial and

$$h = \prod_{i=1}^d t_{\alpha_i} - \prod_{i=1}^d t_{\beta_i} \in J_{P,Q} \text{ such that } \alpha_i(p) = \beta_i(p) \text{ for all } p \in \mathcal{I}'. \tag{3}$$

Since $|\mathcal{I}'| > |\mathcal{I}|$, the desired conclusion follows by induction.

Since

$$\{\varphi_1(p'), \varphi_2(p'), \ldots, \varphi_d(p')\} = \{\psi_1(p'), \psi_2(p'), \ldots, \psi_d(p')\}$$

there exists a permutation π such that $\psi_i(p') = \varphi_{\pi(i)}(p')$. Since we choose $\mathcal{I}$ to be maximal, it follows that $\psi_i(p') \neq \varphi_i(p')$ for some i. Hence, we see that π is nontrivial. Furthermore, if p' has a lower neighbor p'', then $p'' \in \mathcal{I}$. Therefore,

$$\varphi_{\pi(i)}(p') = \psi_i(p') \geq \psi_i(p'') = \varphi_i(p'').$$

Hence, $g = \prod_{i=1}^d t_{\varphi_i} - \prod_{i=1}^d t_{\varphi'_i} \in J_{P,Q}$ is a special where,

$$\varphi'_i(p) = \begin{cases} \varphi_{\pi(i)}(p), & \text{if } p \geq p', \\ \varphi_i(p), & \text{otherwise.} \end{cases} \tag{4}$$

Let $h = \prod_{i=1}^d t_{\varphi'_i} - \prod_{i=1}^d t_{\psi_i}$, then $f = g + h$. Since $f, g \in J_{P,Q}$, it implies that $h \in J_{P,Q}$. Moreover, h satisfies (3). This proves the theorem. □

3. Squarefree binomials

In this section we will prove that $J_{P,Q}$ is generated by squarefree binomials if each connected component of P is rooted or co-rooted poset. In general, without any restriction on P and Q, we have

Remark 3.1. Let P and Q be two posets. Then all quadratic binomials in $J_{P,Q}$ are squarefree. Indeed, assume that $f = t_{\varphi_1}^2 - t_{\psi_1} t_{\psi_2}$. Then $\{\varphi_1(p), \varphi_1(p)\} = \psi_1(p), \psi_2(p)\}$, for all $p \in P$. This implies that $\psi_1 = \psi_2 = \varphi_1$ and hence $f = 0$. This implies that any non-trivial quadratic binomial in $J_{P,Q}$ is squarefree.

We first prove the following

Lemma 3.2. *Any special binomial of type (p, π) can be written as a linear combination of special binomials of type (p, π') where the π' are cyclic permutations.*

Proof. Let $\pi = \pi_1 \pi_2 \dots \pi_l$ be the unique cycle decomposition of π.Then $f = \prod_{i=1}^l f_i^+ - \prod_{i=1}^l f_i^-$ where each $f_i = f_i^+ - f_i^-$ is a special binomial of type (p, π_i). From this presentation of f, it can be shown by induction on l that f can be written as a linear combination of the f_i. □

Now we prove the main theorem.

Theorem 3.3. *Let P be a poset whose connected components are rooted or co-rooted posets. Then for any Q, the ideal $J_{P,Q}$ is generated by squarefree special binomials.*

Proof. By using Theorem 1.1, we may assume that P is connected. Furthermore, we may also assume that P is rooted. We will show by induction on degree of f that f is a linear combination of squarefree binomials in $J_{P,Q}$.

Let $f = t_{\varphi_1} t_{\varphi_2} \cdots t_{\varphi_d} - t_{\psi_1} t_{\psi_2} \cdots t_{\psi_d} \in J_{P,Q}$ and $f \neq 0$. By Theorem 2.2 and the Lemma 3.2, we may assume that f is a special binomial of type (p, π) where π is the cyclic permutation given by $(1\ 2\ \dots\ d)$. By Remark 3.1, we may assume that $d > 2$. If all φ_i as well as all ψ_i are pairwise distinct then f is squarefree and there is nothing to prove. Assume that $\varphi_i = \varphi_j$.

Let $g = g^+ - g^-$ where $g^+ = t_{\varphi_i} t_{\varphi_{i+1}} \dots t_{\varphi_{j-1}}$ and $g^- = t_{\psi_i} t_{\psi_{i+1}} \dots t_{\psi_{j-1}}$. Then $g \in J_{P,Q}$ because it is special of type (p, π') with respect to $\varphi_i, \varphi_{i+1}, \dots, \varphi_{j-1}$ and $\pi' = (i\ i+1\ \cdots\ j-1)$. Indeed, if $p' \not\geq p$ then $\psi_r(p') = \varphi_r(p')$, and if $p' \geq p$ then

$$\psi_r(p') = \varphi_{r+1}(p') \text{ for } r = i, \ldots, j-1 \text{ and } \varphi_j(p') = \varphi_i(p')$$

Hence, we may write

$$f = \frac{f^+}{g^+}\, g + g^-\, h$$

where

$$h = \frac{f^+}{g^+} - \frac{f^-}{g^-}$$

Since $\deg g, \deg h < \deg f$, by induction we know that g and h can be written as a linear combination of squarefree binomials. This completes the proof. □

4. Quadratic binomials

Here we will prove that $J_{P,Q}$ is generated by quadratic squarefree binomials if each connected components of P is rooted or co-rooted and Q does not contain any proper poset cycle of length ≥ 6.

Let Q be a poset. A sequence $q_1, q_2, \ldots, q_{2m}$ of elements in Q is called a *poset cycle* of length $2m$ if $q_1 \leq q_2 \geq q_3 \leq \ldots \leq q_{2m} \geq q_1$.

We say that a poset cycle of length ≥ 6 has a chord if there exist odd i and even j with $j \neq i-1, i+1$ such that $q_i \leq q_j$. (Here 0 is identified with $2m$).

A poset cycle is called *proper* if all elements q_i in the cycle are pairwise distinct and the element with even indexes are pairwise incomparable as well as elements with odd indexes.



Figure 1. A proper poset cycle of length 10

Lemma 4.1. *Let the sequence $\mathcal{C} : q_1, q_2, \ldots, q_{2m}$ of length ≥ 6 be a poset cycle in Q. If $\mathcal{C}$ is not a proper poset cycle then it admits a chord.*

Proof. If $\mathcal{C}$ is not a proper poset cycle then we have one of the following cases:

(i) $q_i \leq q_j$ where both i and j are even or odd.
(ii) $q_i = q_j$ for some odd i and even j.

In case (i), if i and j are even then we have $q_{i-1} \leq q_i \leq q_j$. Then the relation $q_{i-1} \leq q_j$ gives us a chord. Similarly, if i and j are odd then $q_i \leq q_j \leq q_{j+1}$ and $q_i \leq q_{j+1}$ gives us the chord.

Now we discuss the case (ii) and assume first that $i < j$. Then $q_{j-1} \leq q_j = q_i$ where $j-1$ and i are both odd. If $j-1 \neq i$, then we are in case (i). Assume now that $j-1 = i$. Then $q_{j+1} \leq q_{j-1}$, and we are again in case (i).

Similar arguments holds when $i > j$. □

Now we are ready to prove the main Theorem of this section.

Theorem 4.2. *Let P and Q be two posets.*

(a) *If P is an antichain then $J_{P,Q}$ is generated by quadratic squarefree binomials.*
(b) *If P is not an antichain and each connected component of P is rooted or co-rooted, then the following conditions are equivalent:*
(i) *$J_{P,Q}$ is generated by quadratic binomials.*
(ii) *$J_{P,Q}$ is generated by quadratic squarefree binomials.*
(iii) *Each proper poset cycle of Q of length ≥ 6 has a chord.*

Proof. By Theorem 1.1, we may assume that P is connected. Then in case (a), the poset P is just a single element and $K[P,Q]$ is a polynomial ring.

(b) Since P is connected, we may assume that P is rooted.

(i) $\Longleftrightarrow$ (ii): The proof follows from Remark 3.1.

(i) $\Rightarrow$ (iii): Let $\mathcal{C} : q_1, q_2, \ldots, q_{2m}$ be a proper poset cycle in Q of length $2m$ with $m \geq 3$. Let p_1 be the minimal element of the rooted poset P. We define the following maps. For $i = 1, \ldots, m$,

$$\varphi_i(p) = \begin{cases} q_{2i-1}, & p = p_1, \\ q_{2i}, & \text{otherwise.} \end{cases}$$

and

$$\psi_1(p) = \begin{cases} q_1, & p = p_1, \\ q_{2m}, & \text{otherwise.} \end{cases}$$

and for $i = 2, \ldots, m-1$

$$\psi_i(p) = \begin{cases} q_{2i-1}, & p = p_1, \\ q_{2i-2}, & \text{otherwise,} \end{cases}$$

Then $f = t_{\varphi_1} \ldots t_{\varphi_m} - t_{\psi_1} \ldots t_{\psi_m} \in J_{P,Q}$ is a special binomial of degree m and type (p, π) for some $p \in P$ with $p \neq p_1$. After relabeling of indexes, we may assume that $\pi = (1\ m\ m-1\ \ldots\ 2)$ and . Then, since by our assumption $J_{P,Q}$ is generated by quadratic binomials, there exists a nonzero quadratic binomial $g = g^+ - g^- \in J_{P,Q}$ such that $g^+ | f^+$. We may assume that $g = t_{\varphi_i} t_{\varphi_j} - t_{\gamma_1} t_{\gamma_2}$ for some $i < j$ and $\varphi_i(p_1) = \gamma_1(p_1)$ and $\varphi_j(p_1) = \gamma_2(p_1)$. Moreover, we have

$$\{\varphi_i(p), \varphi_j(p)\} = \{\gamma_1(p), \gamma_2(p)\} \text{ for all } p \in P$$

Since $g \neq 0$ there exists $p \in P$ and $p \neq p_1$ such that $\varphi_i(p) = \gamma_2(p)$ and $\varphi_j(p) = \gamma_1(p)$. Therefore, $\varphi_i(p) = q_{2i} \geq \varphi_j(p_1) = q_{2j-1}$ and $\varphi_j(p) = q_{2j} \geq \varphi_i(p) = q_{2i-1}$. The sequence $\mathcal{C}' = q_{2i-1}, q_{2i}, q_{2j-1}, q_{2j}$ is a 4-cycle in Q. In particular $\mathcal{C}$ has a chord.

(iii) $\Rightarrow$ (i): Let $f \in J_{P,Q}$ and $f \neq 0$. Then by Theorem 2.2, we may assume that $f = t_{\varphi_1} \ldots t_{\varphi_m} - t_{\varphi'_1} \ldots t_{\varphi'_m} \in J_{P,Q}$ is a special binomial of type (p, π) where π is a cyclic permutation. After a relabeling of indxes of φ_i and φ'_i, we may assume that $(1\ m\ m-1\ \cdots\ 2)$. If $m = 2$, then the assertion is trivial. Let $m \geq 3$. Since $f \neq 0$ there exists a lower neighbor of p' of p which is unique because P is rooted. Then

the sequence $\mathcal{C} : q_1, q_2, \ldots, q_{2m}$ with $q_{2i-1} = \varphi_i(p')$ and $q_{2i} = \varphi_i(p)$ is a poset cycle in Q.

By Lemma 4.1 and (iii), we know that $\mathcal{C}$ admits a chord. We may assume that the chord is given by the relation $q_1 \leq q_{2i}$ with $i \neq m$.

Let $g = t_{\psi_1} \prod_{j=i+1}^{m} t_{\varphi_j} - t_{\varphi'_1} \prod_{j=i+1}^{m} t_{\varphi'_j}$ where

$$\psi_1(p'') = \begin{cases} \varphi_i(p''), & p'' \geq p, \\ \varphi_1(p''), & \text{otherwise.} \end{cases}$$

By using Lemma 2.1, it can be seen that ψ_1 is isotone. Observe that $g \in J_{P,Q}$. To see this, we have to show that for all $p'' \in P$

$$\{\psi_1(p''), \varphi_{i+1}(p''), \ldots, \varphi_m(p'')\} = \{\varphi'_1(p''), \varphi'_{i+1}(p''), \ldots, \varphi'_m(p'')\}$$

as multi-set.

If $p'' \geq p$, then $\psi_1(p'') = \varphi_i(p'')$ and $\varphi'_j(p'') = \varphi_{j-1}(p'')$ for $j = i+1, \ldots, m$ and $\varphi'_1(p'') = \varphi_m(p'')$. In this case, the two multi-set coincides.

If $p'' \not\geq p$, then $\psi_1(p'') = \varphi_1(p'')$ and $\varphi'_j(p'') = \varphi_j(p'')$ for $j = 1, i+1, \ldots, m$. Again, the two multi-sets coincides.

Then

$$f - t_{\varphi'_2} \ldots t_{\varphi'_i} g = \prod_{j=i+1}^{m} t_{\varphi_j} h,$$

where

$$h = t_{\varphi_1} t_{\varphi_2} \ldots t_{\varphi_i} - t_{\psi_1} t_{\varphi'_2} \ldots t_{\varphi'_i}.$$

Since $f, g \in J_{P,Q}$ and $J_{P,Q}$ is a prime ideal, it follows that $h \in J_{P,Q}$. Note that $\deg g, \deg h < \deg f$. Then by induction, we know that g and h can be written as linear combination of quadratic binomials, which completes the argument.

□

References

[1] M. Bigdeli, J. Herzog, T. Hibi, A. Shikama, A. A. Qureshi, Isotonian algebras.

[2] G. Fløystad, B.M. Greve and J. Herzog, Letterplace and co-letterplace ideals of posets,

[3] V. Ene, J. Herzog and F. Mohammadi, Monomial ideals and toric rings of Hibi type arising from a finite poset, European J. Combin. **32** (2011), 404–421.

[4] J. Herzog and T. Hibi, Monomial ideals, Graduate Texts in Mathematics **260**, Springer, London, 2010.

[5] J. Herzog, A. A. Qureshi and A. Shikama, Alexander duality for monomial ideals associated with isotone maps between posets, to appear in J. Algebra and its Applications.

[6] T. Hibi, Distributive lattices, affine semigroup rings and algebras with straightening laws, Commutative Algebra and Combinatorics (M. Nagata and H. Matsumura, Eds.) Adv. Stud. Pure Math. **11**, North-Holland, Amsterdam, 1987, 93–109.

[7] V. Ene, J. Herzog and S. S. Madani, A note on the regularity of Hibi rings, Manuscripta Math., **148** (2015), 501–506.

[8] M. Juhnke-Kubitzke, L. Katthän and S. S. Madani, Algebraic properties of ideals of poset homomorphisms,

[9] G. Fløystad, A. Nematbakhsh, Rigid ideals by deforming quadratic letterplace ideals,

[10] A. D'Alí, G. Fløystad, A. Nematbakhsh, Resolutions of co-letterplace ideals and generalizations of Bier spheres,
[11] A. D'Alí, G. Fløystad, A. Nematbakhsh, Resolutions of letterplace ideals of posets,

Jürgen Herzog, Fachbereich Mathematik, Universität Duisburg-Essen, Fakultät für Mathematik, 45117 Essen, Germany
E-mail address: juergen.herzog@uni-essen.de

Ayesha Asloob Qureshi, Sabanci University, Orta Mahalle, Tuzla 34956, Istanbul, Turkey
E-mail address: aqureshi@sabanciuniv.edu

Akihiro Shikama, Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Toyonaka, Osaka 560-0043, Japan
E-mail address: a-shikama@cr.math.sci.osaka-u.ac.jp